

Supersymmetric Model of Radiative Seesaw Majorana Neutrino Masses

ERNEST MA

Physics Department, University of California, Riverside, California 92521

ABSTRACT. The radiative seesaw mechanism proposed recently is minimally extended to include supersymmetry in a specific model. Relevant related issues such as leptogenesis and dark matter are discussed.

If the standard model (SM) of particle interactions is extended to include three heavy neutral singlet Majorana fermions N_i ($i = 1, 2, 3$) and a second scalar doublet $\eta = (\eta^+, \eta^0)$ with zero vacuum expectation value (VEV), together with an exactly conserved discrete Z_2 symmetry under which N_i and η are odd and all SM particles are even, then a radiative seesaw mechanism is obtained [1, 2, 3] for small Majorana neutrino masses. The key is a quartic scalar term

$$\lambda_5(\Phi^\dagger\eta)^2 + H.c., \quad (1)$$

where $\Phi = (\phi^+, \phi^0)$ is the SM Higgs doublet, which is allowed under Z_2 . However, such a term is not available in a supersymmetric context. Hence a supersymmetric version of this mechanism is not so straightforward. Nevertheless, it may be accomplished in a minimal extension, as shown below.

Consider the superfields listed in Table 1. Whereas L_i , l_i^c ($i = 1, 2, 3$) are the usual lepton superfields of the minimal supersymmetric standard model (MSSM) and $\Phi_{1,2}$ the usual Higgs superfields, N_i are new Majorana lepton superfields, $\eta_{1,2}$ and χ are new scalar doublet and singlet superfields respectively. The $Z_2 \times Z_2'$ discrete symmetry serves to distinguish the three different kinds of doublets L_i , Φ_1 , η_1 , as well as the

Table 1: Particle content of proposed model.

Superfield	$SU(2) \times U(1)$	Z_2	Z'_2
$L_i = (\nu_i, l_i)$	$(2, -1/2)$	-	+
l_i^c	$(1, 1)$	-	+
$\Phi_1 = (\phi_1^0, \phi_1^-)$	$(2, -1/2)$	+	+
$\Phi_2 = (\phi_2^+, \phi_2^0)$	$(2, 1/2)$	+	+
N_i	$(1, 0)$	-	-
$\eta_1 = (\eta_1^0, \eta_1^-)$	$(2, -1/2)$	+	-
$\eta_2 = (\eta_2^+, \eta_2^0)$	$(2, 1/2)$	+	-
χ	$(1, 0)$	+	-

two different kinds of singlets N_i , χ and doublets Φ_2 , η_2 . Consequently, the superpotential of this model is restricted to be of the form

$$\begin{aligned}
 W = & f_{ij} L_i l_j^c \Phi_1 + h_{ij} L_i N_j \eta_2 + \lambda_1 \Phi_1 \eta_2 \chi + \lambda_2 \Phi_2 \eta_1 \chi \\
 & + \mu_\phi \Phi_1 \Phi_2 + \mu_\eta \eta_1 \eta_2 + \frac{1}{2} \mu_\chi \chi \chi + \frac{1}{2} M_{ij} N_i N_j.
 \end{aligned}
 \tag{2}$$

Since η_2^0 has zero VEV, N_j are not the Dirac mass partners of ν_i and the canonical seesaw mechanism [4] is not operative. However, the Yukawa couplings h_{ij} are available and radiative neutrino masses are obtained [1] with the help of χ , as shown in Figure 1.




Figure 1: One-loop radiative contributions to neutrino mass.

There are two important consequences of this supersymmetric radiative seesaw mechanism. One is leptogenesis [5]. The decay of the

lightest N_i into $L\eta_2$ and their antiparticles will generate a lepton asymmetry which gets converted into the observed baryon asymmetry of the Universe through sphalerons at the electroweak phase transition [6]. On the other hand, the formula for neutrino mass is suppressed by at least a loop factor of $16\pi^2$ as compared to that of the canonical seesaw. This means that the Davidson-Ibarra bound [7] on leptogenesis is reduced by at least two orders of magnitude and the lightest N_i needs to be no heavier than about 10^7 GeV, allowing it to be comfortably below a possible gravitino bound from the reheating of the Universe after inflation [8].

The other is dark matter [9]. As in the MSSM, R parity is conserved in this model. The lightest particle with $R = -1$ is stable and a candidate for the dark matter of the Universe. Similarly, the lightest particle odd under Z'_2 is also stable. In fact, consider the three lightest particles with $(R, Z'_2) = (-, +)$, $(+, -)$, and $(-, -)$ respectively. If one is heavier than the other two combined, then the latter are the two components of dark matter. If not, then all three contribute. In other words, dark matter may not be as boring as usually assumed. It may consist of a rich variety of different stable particles.

Except for N_i , the new particles of this model, i.e. $\eta_{1,2}$ and χ , are expected to be at the TeV scale and should be observable at the forthcoming Large Hadron Collider (LHC).

This work was supported in part by the U. S. Department of Energy under Grant No. DE-FG03-94ER40837.

References

- [1] E. Ma, Phys. Rev. **D73**, 077301 (2006).
- [2] J. Kubo, E. Ma, and D. Suematsu, Phys. Lett. **642**, 18 (2006).
- [3] E. Ma, Mod. Phys. Lett. **A21**, 1777 (2006).
- [4] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Z. Freedman (North-Holland, Amsterdam, 1979), p. 315; T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto (KEK Report No. 79-18, Tsukuba, Japan, 1979), p. 95; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [5] M. Fukugita and T. Yanagida, Phys. Lett. **B174**, 45 (1986).
- [6] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, Phys. Lett. **155B**, 36 (1985).
- [7] S. Davidson and A. Ibarra, Phys. Lett. **B535**, 25 (2002).
- [8] K. Kohri, T. Moroi, and A. Yotsuyanagi, Phys. Rev. **D73**, 123511 (2006).

- [9] For a recent review, see for example G. Bertone, D. Hooper, and J. Silk, Phys. Rept. **405**, 279 (2005).

(Manuscrit reçu le 14 juillet 2006)